



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$F'(y+z) - F'(y) = 0.$$

Then $F'(y)$ must be a constant and $F(y)$ will have the form

$$F(y) = ay + b.$$

Also $f(x)$ and $\theta(x)$ will have the relations, respectively :

$$f(2x) = af(x) + b, \quad \theta(2x) = a\theta(x).$$

Therefore both $f(x)$ and $\theta(x)$ must be constants. Hence we have the following theorem :

The uniform analytical function satisfying a given addition theorem is the only one which satisfies the equation acquired by making $x=y$ on the addition theorem.

DEPARTMENTS.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

The following note will be of value, since it calls attention to the fact that unless the law of a series is given, the *definite* determination of the series is impossible. If only a certain number of terms of a series is given and not the law of the series, the actual law of the series can in no case be really determined. All that can be done is to find the *simplest law* which the few given terms will obey. The solutions referred to may be justified on the ground that they were attempts to find this *simplest law*. ED. F.

NOTE ON THE SOLUTION OF PROBLEM 266, IN AUGUST-SEPTEMBER MONTHLY.

By CLARENCE E. COMSTOCK, Bradley Polytechnic Institute.

Find the n th term and the sum of n terms of the series $1+3+7+17+\dots$

One writer assumes without the least excuse for such an assumption that it is an arithmetical series of the third order and proceeds by the method of finite differences, getting a result in accord with his assumption. Five terms of his series are $1+3+7+17+37+\dots$

Another writer assumes with a little more plausible excuse that it is a recurring series of the second order and, of course, gets a result in accord with his assumption. Five terms of his series are $1+3+7+17+41+\dots$

Suppose I make the assumption that it is an arithmetical series of the fourth order. I can then build up

$$1+3+7+17+38+76+\dots \text{ or } 1+3+7+17+39+81+\dots$$

or as many more as I care to take the time to construct, on the one supposition that it is an arithmetical series of the fourth order. Or let me assume it a recurring series of the fourth order, and get, using the scale of relation,

$$u_n = 2u_{n-1} + u_{n-2} + 0u_{n-3} + u_{n-4}, \text{ say, } 1+3+7+17+42+104+\dots$$

The fact is, the problem is absolutely without meaning. There is nothing in the sequence $1+3+7+17+\dots$ to show what the next term is. A series is not determined until its law is in some way stated or indicated.

267 Proposed by O. E. GLENN, Ph. D., Philadelphia, Pa.

Express the trigonometric functions of x as infinite continued fractions.

Solution by R. D. CARMICHAEL, Professor of Mathematics, Anniston, Ala.; and J. SCHEFFER, A. M., Kee Mar College, Hagerstown, Md.

$$\text{Let the series } u = A - B + C - D \dots = \frac{a_1}{c_1} + \frac{a_2}{c_2} + \frac{a_3}{c_3} + \dots$$

To determine the a 's we have the following (Euler's "Introductio," I, §367):

$$a_1 = A.c_1, a_2 = \frac{B.c_1c_2}{A-B}, a_3 = \frac{AC.c_2c_3}{(A-B)(B-C)}, a_4 = \frac{BD.c_3c_4}{(B-C)(C-D)} \dots$$

in which the c 's are to be so chosen that the a 's are integral functions. Applying these formulas, we have

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots = \frac{x}{1} + \frac{x^2}{2.3-x^2} + \frac{2.3x^2}{4.5-x^2} + \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots = \frac{1}{1} + \frac{x^2}{1.2-x^2} + \frac{1.2x^2}{3.4-x^2} + \frac{3.4x^2}{5.6-x^2} + \dots$$

$$\begin{aligned} \tan x &= \frac{2}{\pi-2x} - \frac{2}{\pi+2x} + \frac{2}{3\pi-2x} - \frac{2}{3\pi+2x} + \frac{2}{5\pi-2x} + \dots^* \\ &= \frac{2}{\pi-2x} + \frac{4x}{(\pi-2x)^2} + \frac{2\pi}{\pi^2-4x^2} + \frac{(\pi-2x)(3\pi-2x)^2}{2x(\pi+2x)} + \dots \end{aligned}$$

**Encyclopedia Britannica*, Vol. XXIII, p. 572.